arXiv:1211.3558v1 [nucl-ex] 15 Nov 2012

Consistency of ^{8}B neutrino spectra

O. S. Kirsebom,[1,*] H. O. U. Fynbo,[2] R. Raabe,[3] K. Riisager,[2] and T. Roger[4]
[1] *TRIUMF, Vancouver, British Columbia, V6T 2A3, Canada*
[2] *Department of Physics and Astronomy, Aarhus University, DK-8000 Aarhus C, Denmark*
[3] *Instituut voor Kern- en Stralingsfysica, KU Leuven, B-3001 Leuven, Belgium*
[4] *GANIL, Bd Henri Becquerel, BP 55027, F-14076 Caen Cedex 05, France*

We identify and quantify systematic effects not accounted for in two previous measurements of the α-α relative-energy distribution in the β decay of ^{8}B, which can explain the apparent disagreement with respect to two newer measurements. This settles a current dispute concerning the shape of the ^{8}B neutrino spectrum of importance to solar-neutrino studies.

PACS numbers: 26.65.+t, 29.30.Ep, 29.40.Wk

Introduction. New solar-neutrino observatories are allowing us to test our understanding of the Sun and of neutrinos themselves with increasing precision. Neutrinos emitted in the β decay of ^{8}B play a central role in these endeavours because they dominate the solar-neutrino spectrum above approximately 2 MeV. Precise and accurate knowledge of the laboratory ^{8}B neutrino spectrum is necessary to detect distortions of the solar-neutrino spectrum [1].

The β decay of ^{8}B populates a broad distribution of excitation energies in ^{8}Be. Subsequently, the excited ^{8}Be nucleus breaks up into two α particles. The ^{8}B neutrino spectrum can be deduced from the α-α relative-energy distribution, which can be determined in a laboratory experiment. Our collaboration has used two different experimental techniques to determine the α-α relative-energy distribution [2, 3]. The two techniques give consistent distributions, which disagree slightly with the distributions determined in two previous experiments by Bhattacharya *et al.* [4] and Winter *et al.* [5, 6]. In fact, the shapes of the former and the latest distributions are in reasonable agreement, but the maxima of the distributions deviate by approximately 20 keV, which results in a 4% difference on the ^{8}B neutrino spectrum at high energies ($E_\nu > 14$ MeV). This uncertainty complicates the attempts being made to detect neutrinos from the fusion of ^{3}He+p in the Sun, and the attempts being made to detect the diffuse supernova neutrino background [7].

The purpose of the present Rapid Communication is to identify and quantify systematic effects not accounted for in the two former measurements of the α-α relative-energy distribution, which can explain the 20 keV deviation relative to the two latest measurements performed by our collaboration, thus settling the current dispute [8, 9].

Response of Si detectors to different ions. Both the former and the latest measurements were made with Si detectors, and in all four cases the energy calibration of these detectors was the main source of uncertainty. Typically, the calibration of a Si detector consists in establishing a linear relationship between the observed pulse height, x, and the deposited energy, E. However, the physical quantity that is directly related to the observed pulse height is not the deposited energy, but the number of collected electron-hole pairs, N. Therefore, it is more correct to assume a linear relationship between x and N, *i.e.*[1]

$$x = aN + b\,. \tag{1}$$

The energy deposited in the detector by a charged particle can be divided into two parts: the part that goes into electron-hole pair formation (ionizing), which we shall denote by E_i, and the part that does not (non-ionizing), which we shall denote by E_n, *i.e.* $E = E_\mathrm{i} + E_\mathrm{n}$. Denoting the electron-hole pair collection efficiency by η and the energy required to create an electron-hole pair by ϵ, we have

$$N = \frac{\eta\, E_\mathrm{i}}{\epsilon} = \frac{\eta}{\epsilon}\,(E - E_\mathrm{n})\,.$$

Introducing $\kappa = \eta/\epsilon$, this simplifies to

$$N = \kappa\,(E - E_\mathrm{n})\,. \tag{2}$$

Finally, combining Eqs. (1) and (2), we find

$$x = a\kappa\,(E - E_\mathrm{n}) + b\,. \tag{3}$$

Note that E_n depends on the charge and the mass of the particle and is a function of the deposited energy, E. Ideally, the parameter κ would be independent of the charge and the mass and would be constant with E. However, the experimental study of Lennard *et al.* [10] shows that κ has a weak dependence on these quantities. (The interpretation of this effect suggested by Lennard *et al.* is that ϵ decreases with increasing stopping power, but this is irrelevant to the present analysis.) Finally, the calibration coefficients a and b are strictly independent of charge, mass, and E; they are determined by the experimental apparatus, *i.e.* the detector and the associated electronics.

---

*Corresponding author: oliverk@triumf.ca

[1] To simplify the discussion, we neglect non-linearities in the electronic processing of the signal (amplification, shaping, and digital conversion).

The experiment of Winter et al. In the experiment of Winter *et al.* [5, 6], the ^{8}B ions were implanted in a Si detector and the linearly added pulse heights of the two α particles were measured. The α particles have nearly equal energies, *i.e.* $E_{\alpha_1} = E_{\alpha_2} = \frac{1}{2}E$. The number of collected electron-hole pairs created by each α particle is given by Eq. (2). The total number of collected electron-hole pairs is simply twice this number, *i.e.*

$$N_{^8\mathrm{B}} = \kappa_\alpha \left[E - 2E_{\mathrm{n},\alpha}(\tfrac{1}{2}E) \right] . \tag{4}$$

The energy calibration was performed by implanting the β-delayed α emitter ^{20}Na in the detector. The linearly added pulse heights of the α particle and the ^{16}O ion were measured. In this case, the energy is shared approximately in ratios of 4:1, *i.e.* $E_\alpha = \frac{4}{5}E$ and $E_{^{16}\mathrm{O}} = \frac{1}{5}E$, and the total number of collected electron-hole pairs is given by

$$\begin{aligned} N_{^{20}\mathrm{Na}} &= \kappa_\alpha \left[\tfrac{4}{5}E - E_{\mathrm{n},\alpha}(\tfrac{4}{5}E) \right] \\ &\quad + \kappa_{^{16}\mathrm{O}} \left[\tfrac{1}{5}E - E_{\mathrm{n},^{16}\mathrm{O}}(\tfrac{1}{5}E) \right] . \end{aligned} \tag{5}$$

In general, $N_{^8\mathrm{B}} \neq N_{^{20}\mathrm{Na}}$, *i.e.* for the same deposited energy, E, different numbers of electron-hole pairs are collected and hence different pulse heights are measured. This is the basic reason why a calibration based on ^{20}Na data cannot be applied to ^{8}B data without proper correction. Note that Eqs. (4) and (5) do not include the small, but significant, contribution of the β particles, which we are not concerned with here. It follows from Eqs. (4) and (5) that the necessary correction for the different response of Si detectors to α particles and ^{16}O ions, is

$$\begin{aligned} \Delta E &= \kappa_\alpha^{-1} \left(N_{^8\mathrm{B}} - N_{^{20}\mathrm{Na}} \right) \\ &= \tfrac{1}{5}E - \frac{\kappa_{^{16}\mathrm{O}}}{\kappa_\alpha} \left[\tfrac{1}{5}E - E_{\mathrm{n},^{16}\mathrm{O}}(\tfrac{1}{5}E) \right] \\ &\quad + E_{\mathrm{n},\alpha}(\tfrac{4}{5}E) - 2E_{\mathrm{n},\alpha}(\tfrac{1}{2}E) . \end{aligned} \tag{6}$$

We will now determine the magnitude of ΔE using the experimental data of Lennard *et al.* [10]. In this experiment, the response of a Si detector to ^{1}H, ^{4}He, ^{7}Li and ^{16}O ions was studied. The energy of the incident ion (E_K in the notation of Lennard *et al.*) was determined by the time-of-flight technique. The energy loss in the detector window (ΔE_w in the notation of Lennard *et al.*) was determined from the rate of change of pulse height with angle. The energy deposited in the detector could then be determined as $E = E_\mathrm{K} - \Delta E_\mathrm{w}$. Finally, the non-ionizing energy deposition (E_n in the present notation and ΔE_n in the notation of Lennard *et al.*) was estimated from theory. First, we use the α-particle data of Lennard *et al.* to calibrate the Si detector used in their experiment, *i.e.* we determine the parameters $a\kappa_\alpha$ and b in Eq. (3). Corresponding values of E_α (deposited energy) and x_α (pulse height) were determined experimentally at 13 energies between 0.53 and 2.96 MeV. A linear fit to the 13 data points ($E_\alpha - E_{\mathrm{n},\alpha}$, x_α) gives[2]



FIG. 1: Reanalysis of the data of Lennard *et al.* [10]. $E_{^{16}\mathrm{O}}$ is the energy deposited in the detector and $\delta E_{^{16}\mathrm{O}}$ is defined in Eq. (7). The dashed line shows the value adopted by us for $\delta E_{^{16}\mathrm{O}}$ in the energy range 0.5–1.2 MeV.

$$a\kappa_\alpha = 0.1843(2)\ \mathrm{keV}^{-1} , \quad b = 0.2(3) ,$$

with $\chi^2/\mathrm{dof} = 1.3$. Next, we consider the ^{16}O data. Corresponding values of $E_{^{16}\mathrm{O}}$ and $x_{^{16}\mathrm{O}}$ were determined experimentally at four energies between 0.73 and 2.52 MeV. Using these data and the values of $a\kappa_\alpha$ and b determined above, we evaluate the following quantity,

$$\delta E_{^{16}\mathrm{O}} = E_{^{16}\mathrm{O}} - \frac{x_{^{16}\mathrm{O}} - b}{a\kappa_\alpha} , \tag{7}$$

which is shown in Fig. 1. We conclude that $\delta E_{^{16}\mathrm{O}}$ is approximately constant in the energy range of interest with regard to the ^{20}Na calibration, $E_{^{16}\mathrm{O}} = 0.5$–$1.2$ MeV, and in this range $\delta E_{^{16}\mathrm{O}} = 66(4)$ keV. Substituting Eq. (3) for $x_{^{16}\mathrm{O}}$ in Eq. (7), we obtain

$$\delta E_{^{16}\mathrm{O}} = E_{^{16}\mathrm{O}} - \frac{\kappa_{^{16}\mathrm{O}}}{\kappa_\alpha} \left[E_{^{16}\mathrm{O}} - E_{\mathrm{n},^{16}\mathrm{O}}(E_{^{16}\mathrm{O}}) \right] ,$$

which we recognize as the first two terms in Eq. (6) with $\frac{1}{5}E = E_{^{16}\mathrm{O}}$. Interpolating the theoretical calculations of Lennard *et al.*, we determine the two last terms of Eq. (6) to be $E_{\mathrm{n},\alpha}(\frac{1}{2}E) = 9.1$ keV and $E_{\mathrm{n},\alpha}(\frac{4}{5}E) = 9.8$ keV with negligible uncertainties [10], for a total energy of $E = 3.0$ MeV corresponding to the maximum of the α-α relative-energy distribution in the β decay of ^{8}B. We thus

---

[2] The data is consistent with an energy-independent κ_α. In contrast, we make no assumptions about $\kappa_{^{16}\mathrm{O}}$.

TABLE I: Maxima of the ^{8}Be excitation-energy distribution obtained in the four most recent studies of the β decay of ^{8}B. The ^{8}Be excitation energy, E_x, and the α-α relative energy, $E_{\alpha\text{-}\alpha}$, are related as $E_x = E_{\alpha\text{-}\alpha} - E_{\mathrm{gs}}$, where $E_{\mathrm{gs}} = 92$ keV is the energy of the ground state of ^{8}Be relative to the 2α threshold. The deviation is computed with respect to the average of Refs. [2] and [3].

Study	Maximum (keV)	1σ uncertainty (keV)	Deviation (keV)
Winter *et al.* [5]	2 943	9	21
Bhattacharya *et al.* [4]	2 939	5	17
Kirsebom *et al.* [2]	2 921	5	−1
Roger *et al.* [3]	2 923^a	6	1

^{a}This is the updated value given in Ref. [9].

conclude that

$$\begin{aligned}\Delta E &= \delta E_{^{16}\mathrm{O}} + E_{\mathrm{n},\alpha}(\tfrac{4}{5}E) - 2E_{\mathrm{n},\alpha}(\tfrac{1}{2}E) \\ &= 66(4)\ \mathrm{keV} + 9.8\ \mathrm{keV} - 2\times 9.1\ \mathrm{keV} \\ &= 58(4)\ \mathrm{keV}\,. \end{aligned} \tag{8}$$

This correction is approximately 13 keV larger than the correction applied by Winter *et al.*, which is said to be "40–50 keV for the various alpha lines, with an uncertainty of 5 keV" [5, 6], citing Lennard *et al.* [10]. The sign and the magnitude of this difference is such that it can account for the 20 keV shift of the distribution of Winter *et al.* relative to our distributions, cf. Table I. We thus conclude that a misinterpretation of the data of Lennard *et al.* on the part of Winter *et al.* is the likely reason for the discrepancy between their result and ours.

^{20}Na was used for energy calibration in both of our experiments, but only the implantation technique [3] is sensitive to the different response of Si detectors to α particles and ^{16}O ions. The internal consistency of our experiments may thus been seen as a verification of Eq. (8).

Winter *et al.* also used an external ^{228}Th α source to provide calibration points at higher energies (5–9 MeV). Corrections were made for the energy loss of the α particles in the source, but it is unclear whether the different implantation depths of the α-unstable daughter nuclei in the ^{228}Th decay series were taken into account. As discussed in Section III of Ref. [2], neglect of this effect will distort the energy calibration.

Reanalysis of ^{12}B, ^{12}N, ^{20}Na decay data. To further support our case, we have reanalyzed data from an experiment conducted by our collaboration where ^{12}B, ^{12}N, and ^{20}Na ions were implanted in a finely segmented Si detector and the linearly added pulse heights of the charged decay products were measured [11]. The same detector was later used in one of our ^{8}B experiments [3]. The ^{12}B, ^{12}N, and ^{20}Na energy spectra are shown in Figs. 2 (a) and (b). The two most prominent peaks in the ^{20}Na spectrum have been used to calibrate the energy axis, taking into account the additional energy deposited by the β particles, as determined with a GEANT4 simulation [3], and taking into account the β-decay recoil energy, which is relatively small. The β decays of ^{12}B and ^{12}N populate excited states in ^{12}C, which breakup into three α particles. For the purpose of estimating the non-ionizing energy loss, $E_{\mathrm{n},\alpha}$, we assume that each α particle gets one-third of the breakup energy. This is a reasonable assumption for the two states in ^{12}C under consideration. Since $E_{\mathrm{n},\alpha}$ only varies slowly with energy, about 0.9 keV per MeV, the error introduced by this approximation is, in any case, very small. In analogy to Eq. (8), we can then write down an expression for the error that results if the ^{20}Na calibration is applied to the ^{12}B and ^{12}N data



FIG. 2: (Color online) Decay energy-spectra of ^{20}Na ions (a) and ^{12}B and ^{12}N ions (b) implanted in a finely segmented Si detector [11]. The peaks are labeled by the excitation energy in the daughter nucleus in MeV.

without correction for the different response of Si detectors to α particles and ^{16}O ions,

$$\Delta E = 66(4)\ \mathrm{keV} + E_{\mathrm{n},\alpha}(\tfrac{4}{5}E) - 3E_{\mathrm{n},\alpha}(\tfrac{1}{3}E)\,. \qquad (9)$$

The data analysis is summarized in Table II. As seen in Fig. 2 (b), the β-decays of ^{12}B and ^{12}N populate two narrow states in ^{12}C, the 0^+ state at 7.654 MeV and the 1^+ state at 12.710 MeV. When the additional energy deposited by the β particles and the β-decay recoil energy are taken into account, we find that the peaks are located above their nominal energies by 54(4) keV and 49(9) keV for ^{12}B and by 56(4) keV and 61(7) keV for ^{12}N. These numbers are in good agreement with the deviations expected based on Eq. (9), 53(4) keV and 48(4) keV, thus lending further support to our case.

The experiment of Bhattacharya et al. In the experiment of Bhattacharya *et al.* [4], the ^{8}B ions were implanted in a thin carbon foil viewed by an arrangement of Si detectors. Before the α particles can deposit their energy in the sensitive volume of the detectors, they must exit the foil and traverse the dead layer of the detector, which leads to a small, but significant, energy loss. Interactions with electrons are frequent and lead to an approximately Gaussian energy-loss distribution, assuming that the layers traversed are not too thin, nor too thick [14, 15]. Interactions with nuclei (Coulomb scattering) are far less frequent and do not contribute to the mean energy loss. However, because individual scattering events in this case can result in a large energy loss, the energy-loss distribution acquires a pronounced *high*-energy tail, which is seen as *low*-energy tail in the response function of the detector. The magnitude of the tail is very sensitive to the thickness traversed [16]. In one of the two ^{8}B experiments performed by our collaboration [2], it was indeed found that the response function determined with a ^{241}Am α source had a more pronounced low-energy tail than the response function determined with ^{20}Na implanted in a thin carbon foil.

The effect of the low-energy tail is to shift the *apparent* position of the maximum of the α-particle energy spectrum toward lower energies. Bhattacharya *et al.* [4] used a ^{148}Gd α source to determine the response function of their detectors, which could have led them to overestimate the magnitude of the low-energy tail. This, in turn, would have led them to conclude that the *actual* position of the maximum of the α-particle energy spectrum is higher in energy than it really is. If the difference were the same as in our experiment [2], the error would be about 8 keV, leading to a $2 \times 8 = 16$ keV error on the α-α relative energy. The sign and the magnitude of this error is such that it can account for the 20 keV shift relative to our distributions. In our experiment, the response function was extracted from the ^{20}Na data. Since the ^{20}Na ions and the ^{8}B ions were implanted in the same carbon foil with similar depth profiles, the low-energy tail should be very similar in the two cases.

Finally, we note that Bhattacharya *et al.* determine the peak of the α-α relative-energy distribution to be slightly wider than we do and Winter *et al.* do, suggesting that experimental broadening effects were not completely accounted for in their experiment.

Summary and conclusion. Our collaboration has determined the α-α relative-energy distribution in the β decay of ^{8}B using two different experimental techniques, which yield consistent distributions [2, 3] that, however, deviate by a small, but significant, amount with respect to two previous determinations by Bhattacharya *et al.* [4] and Winter *et al.* [5, 6]. In the present Rapid Communication, we have identified possible sources of error in the energy calibrations of the previous experiments that may account for the discrepancy. We suggest that Winter *et al.* did not correctly account for the different response of Si detectors to α particles and ^{16}O ions. We have presented empirical evidence supporting our suggestion and estimated the associated error to be approximately +13 keV. We suspect that Bhattacharya *et al.* may have overestimated the magnitude of the low-energy tail of their response function and we have argued that the associated error could be as large as +16 keV. As an important corollary, the ^{8}B neutrino spectra deduced from the four experiments now agree within about 1% at high neutrino energies, whereas before they differed by 4%.

---

[1] W. C. Haxton, R. G. H. Robertson, and A. M. Serenelli, arXiv:1208.5723v1 [astro-ph.SR] .

[2] O. S. Kirsebom *et al.*, Physical Review C **83**, 065802 (2011).

[3] T. Roger *et al.*, Physical Review Letters **108**, 162502 (2012).

[4] M. Bhattacharya, E. G. Adelberger, and H. E. Swanson, Physical Review C **73**, 055802 (2006).

[5] W. T. Winter *et al.*, Physical Review Letters **91**, 252501 (2003).

[6] W. T. Winter, S. J. Freedman, K. E. Rehm, and J. P. Shiffer, Physical Review C **73**, 025503 (2006).

[7] B. Aharmim *et al.*, The Astrophysical Journal **653**, 1545 (2006).

[8] S. J. Freedman, K. E. Rehm, J. P. Schiffer, and D. Seweryniak, Phys. Rev. Lett. **109**, 189201 (2012).

[9] T. Roger, O. S. Kirsebom, H. O. U. Fynbo, and R. Raabe, Phys. Rev. Lett. **109**, 189202 (2012).

[10] W. N. Lennard *et al.*, Nuclear Instruments and Methods in Physics Research A **248**, 454 (1986).

[11] S. Hyldegaard *et al.*, Physics Letters B **678**, 459 (2009).

[12] D. R. Tilley *et al.*, Nuclear Physics A **636**, 249 (1998).

[13] F. Ajzenberg-Selove, Nuclear Physics A **506**, 1 (1990).

[14] L. Glazov, Nuclear Instruments and Methods in Physics Research B **161-163**, 1 (2000).

[15] L. G. Glazov, Nuclear Instruments and Methods in Physics Research B **192**, 239 (2002).

[16] L. Buchmann, G. Ruprecht, and C. Ruiz, Physical Re-

TABLE II: Reanalysis of the experimental data of Ref. [11]. E_x is the excitation energy of the daughter nucleus and E is the breakup energy, calculated as $E = E_x - S_\alpha$ for ^{20}Na and as $E = E_x - S_{3\alpha}$ for ^{12}B and ^{12}N, where $S_\alpha = 4\,729.8$ keV is the $\alpha + {}^{16}$O threshold energy in ^{20}Ne and $S_{3\alpha} = 7\,274.7$ keV is the 3α threshold energy in ^{12}C; both are known to high precision. E_β is the shift due to the energy deposited by the β particles, determined as the mean of the energy-loss distribution obtained with a GEANT4 simulation [3]. E_{R} is the β-decay recoil energy averaged over lepton angles, cf. Section IV in Ref. [2]. $E' = E + E_\beta + E_{\mathrm{R}}$ is the breakup energy corrected for the β- and recoil shifts. E_{exp} is the mean energy of the peaks in Figs. 2 (a) and (b). The ^{20}Na peaks were used to calibrate the energy axis. $\Delta E_{\mathrm{exp}} = E_{\mathrm{exp}} - E'$ is the deviation with respect to the corrected breakup energy. Finally, ΔE is the expected deviation based on Eq. (9). All energies are in keV.

	E_x [12, 13]	E	E_β	E_{R}	E'	E_{exp}	ΔE_{exp}	ΔE
^{20}Na	7 421.9(12)	2 692.1(12)	26(2)	0.5	2 719(2)	...	...	...
	10 273.2(19)	5 543.4(19)	31(2)	0.2	5 575(3)	...	...	...
^{12}B/^{12}N	7 654.2(15)	379.5(15)	28(2)/26(2)	1.2/2.7	409(3)/408(3)	462(3)/464(3)	54(4)/56(4)	53(4)[a]
	12 710(6)	5 435(6)	59(2)/30(2)	0.0/0.5	5 494(6)/5 466(6)	5 543(6)/5 527(3)	49(9)/61(7)	48(4)

[a] An additional correction of +1.3 keV was necessary to account for the energy dependence of $E_{\mathrm{n},\alpha}$ outside the energy range covered by the ^{20}Na calibration data.

view C **80**, 045803 (2009).